

STT102 List of eExam Questions in the Bank

Latex formatted questions may not properly render

Q1 In the age distribution of Recipients of Nursing scholarship of 25, 21, 22, 20, 19, 30, 27, 28, 32 and 18. The variance is _____. (Hint: use $S^2 = \sum (X - \bar{X})^2$)

Q2 If $X = 10, 12, 8, 7, 5$. $\sum_{i=1}^5 X_i$ is

Q3 Let $Y = 2, 5, 6, 7$. $\sum_{j=1}^4 Y_j$ has the value

Q4 Take $X = 29, 27, 28, 30, 35$. $\bar{X}$ is

Q5 One Precaution in correlation is that

Q6 The scores obtained by 10 students in a practical class are as follows: 20, 50, 30, 40, 60. The mean score is

Q7 In the distribution having classes 0-4 5-9 10-14 15-19 20-24 . . . The upper class boundary for class 3 is

Q8 Mean, Median and Mode are measures of

Q9 In attitude test, the scores for 5 newly students are as stated here, Attitude: 5, 4, 3, 2, 1. The percentage attributable to attitude of score 3 is

Q10 Cluster Sampling is one whose members are

Q11 Systematic Sampling is random sampling method

Q12 Simple Random Sampling (SRS) is one for which each possible sample is likely to be selected

Q13 The sample characteristics is

Q14 is An example of population characteristics

Q15 Statistics is the that deals with data collection, and summarising facts which are expressible in numerical form

Q16 The following data were collected on ten infants. Find the standard error, $\sqrt{S_{yx}}$. Where $S_{yx}^2 = \sum_{i=1}^{10} (y_i - \hat{y}_i)^2$ and y_i are the observed values, $\hat{y}_i$ are the predicted values

$\sqrt{S_{yx}} = 5.75$

$\sqrt{S_{yx}} = 4.75$

$\sqrt{S_{yx}} = 2.75$

$\sqrt{S_{yx}} = 3.75$

Q17 Given the general form of linear equation $y = b + b_1X$. If $b_1 > 0$, then the line slopes

downward
upward
flat
parallel

Q18 Consider Attitude Scores for five newly admitted Nursing students towards alcoholic patients below: Attitude: 5, 4, 3, 2, 1 . The percentage due to attitude 3 is _____

- 0.5
- 0.4
- 0.2
- 0.3

Q19 The data below represent systolic blood pressure readings (mm Hg), using Spearman's Rank Order Correlation method, determine correlation coefficient $\sqrt{r_{\{s\}}}$ of the two readings.

- $\sqrt{r_{\{s\}}} = 0.23$
- $\sqrt{r_{\{s\}}} = -0.32$
- $\sqrt{r_{\{s\}}} = 0.32$
- $\sqrt{r_{\{s\}}} = -0.23$

Q20 Determine Correlation Coefficient 'r' using the above values or from your direct-calculation

- 0.9
- 0.91
- 0.95
- 0.92

Q21 Find the value of $\sqrt{S_{\{w_1w_2\}}}$ in question one above

- 4135
- 4235
- 4335
- 4325

Q22 From the above, evaluate $\sqrt{S_{\{w_2w_2\}}}$.

- 2440
- 2410
- 2420
- 2430

Q23 This is for Questions 1 to 4. Two weekly scores of a students are as below <> . Find $\sqrt{S_{\{w_1w_1\}}}$

- 6250.25
- 6150.5
- 6312.5
- 6300.5

Q24 Given that $X = 20, 30, 40, 50, 60$. Find $\sqrt{\bar{X}}$.

- 40,

30,
35
45

Q25 Consider this distribution 12, 20, 13, 15, 17, 15, 18. Find $\bar{X}_{\{m\}}$, where $\bar{X}_{\{m\}}$ is as earlier defined.

9
11
13
15

Q26 Let $\bar{X}_{\{m\}}$ be the Median Score, Determine $\bar{X}_{\{m\}}$ in 15, 13, 15, 12, 12, 16, 15, 14, 13

10
12
14
16

Q27 Suppose $X_{\{m\}}$ is the Mode. Find $X_{\{m\}}$ in 15, 13, 15, 12, 12, 16, 15, 14, 13.

11
13
15
17

Q28 Suppose $X = 10, 12, 8, 7, 5$. Find the value of $\sum_{i=1}^5 X_{\{i\}}^2$

204
214
224
234

Q29 Determine $\sum_{i=1}^5 X_{\{i\}}^2$ if $X = 10, 12, 8, 7, 5$

1265
1764
1785
1951

Q30 Let $Y = 2, 5, 6, 7$. Find $\sum_{j=1}^4 Y_{\{j\}}$

114
120
125
141

Q31 If $X=10, 12, 8, 7, 5$ Determine $\sum_{i=1}^5 X_{\{i\}}$

40
41
42
43

Q32 Given that $X = 20, 30, 40, 50, 60$. Find $\bar{X}$.

- 40,
- 30,
- 35
- 45

Q33 Consider this distribution 12, 20, 13, 15, 17, 15, 18. Find $\bar{X}_{\{m\}}$, where $\bar{X}_{\{m\}}$ is as earlier defined.

- 9
- 11
- 13
- 15

Q34 Let $\bar{X}_{\{m\}}$ be the Median Score, Determine $\bar{X}_{\{m\}}$ in 15, 13, 15, 12, 12, 16, 15, 14, 13

- 10
- 12
- 14
- 16

Q35 Suppose $\bar{X}_{\{m\}}$ is the Mode. Find $\bar{X}_{\{m\}}$ in 15, 13, 15, 12, 12, 16, 15, 14, 13

- 11
- 13
- 15
- 17

Downloaded from noughgeeks.com